

## Medication Guide

### **DURAGESIC® (Dur-ah-GEE-zik) (fentanyl) Transdermal System, CII**

#### **DURAGESIC® is:**

- A strong prescription pain medicine that contains an opioid (narcotic) that is used to manage pain severe enough to require daily around-the-clock, long-term treatment with an opioid, in people who are already regularly using opioid pain medicine, when other pain treatments such as non-opioid pain medicines or immediate-release opioid medicines do not treat your pain well enough or you cannot tolerate them.
- A long-acting (extended-release) opioid pain medicine that can put you at risk for overdose and death. Even if you take your dose correctly as prescribed you are at risk for opioid addiction, abuse, and misuse that can lead to death.
- Not for use to treat pain that is not around-the-clock.

#### **Important information about DURAGESIC®:**

- **Get emergency help right away if you use too much DURAGESIC® (overdose).** When you first start taking DURAGESIC®, when your dose is changed, or if you take too much (overdose), serious or life threatening breathing problems that can lead to death may occur.
- Never give anyone else your DURAGESIC®. They could die from using it. Store DURAGESIC® away from children and in a safe place to prevent stealing or abuse. Selling or giving away DURAGESIC® is against the law.
- If the patch accidentally sticks to a family member while in close contact, take the patch off, wash the area with water, and get emergency help right away because an accidental exposure to DURAGESIC® can lead to death or other serious medical problems.
- Proper disposal of DURAGESIC® after use and for unused patches when no longer needed: Fold the sticky sides of the patch together and flush down the toilet. Do not put patches in a trash can.

#### **Do not use DURAGESIC® if you have:**

- severe asthma, trouble breathing, or other lung problems.
- a bowel blockage or have narrowing of the stomach or intestines.

#### **Before applying DURAGESIC®, tell your healthcare provider if you have a history of:**

- head injury, seizures
- liver, kidney, thyroid problems
- problems urinating
- pancreas or gallbladder problems
- abuse of street or prescription drugs, alcohol addiction, or mental health problems.

#### **Tell your healthcare provider if you:**

- have a fever
- **are pregnant or planning to become pregnant.** Prolonged use of DURAGESIC® during pregnancy can cause withdrawal symptoms in your newborn baby that could be life-threatening if not recognized and treated.
- **are breastfeeding.** DURAGESIC® passes into breast milk and may harm your baby.
- are taking prescription or over-the-counter medicines, vitamins, or herbal supplements. Taking DURAGESIC® with certain other medicines can cause serious side effects that could lead to death.

#### **When using DURAGESIC®:**

- Do not change your dose. Apply DURAGESIC® exactly as prescribed by your healthcare provider.
- See the detailed Instructions for Use for information about how to apply and dispose of the DURAGESIC® patch.
- Do not apply more than 1 patch at the same time unless your healthcare provider tells you to.
- You should wear the DURAGESIC® patch continuously for 3 days, unless advised otherwise by your healthcare provider.
- **Call your healthcare provider if the dose you are using does not control your pain.**
- **Do not stop using DURAGESIC® without talking to your healthcare provider.**

#### **While using DURAGESIC® DO NOT:**

- Take hot baths or sunbathe, use hot tubs, saunas, heating pads, electric blankets, heated waterbeds, or tanning lamps, or engage in exercise that increases your body temperature. These can cause an overdose that can lead to death.
- Drive or operate heavy machinery, until you know how DURAGESIC® affects you. DURAGESIC® can make you sleepy, dizzy, or lightheaded.
- Drink alcohol or use prescription or over-the-counter medicines that contain alcohol. Using products containing alcohol during treatment with DURAGESIC® may cause you to overdose and die.

#### **The possible side effects of DURAGESIC® are:**

- constipation, nausea, sleepiness, vomiting, tiredness, headache, dizziness, abdominal pain, itching, redness, or rash where the patch is applied. Call your healthcare provider if you have any of these symptoms and they are severe.

#### **Get emergency medical help if you have:**

- trouble breathing, shortness of breath, fast heartbeat, chest pain, swelling of your face, tongue or throat, extreme drowsiness, light-headedness when changing positions, or you are feeling faint.

These are not all the possible side effects of DURAGESIC®. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. **For more information go to [dailymed.nlm.nih.gov](http://dailymed.nlm.nih.gov)**

Manufactured by: Alza Corporation Vacaville CA 95688; Manufactured for: Janssen Pharmaceuticals Inc. Titusville NJ 08560 [www.Duragesic.com](http://www.Duragesic.com) or call 1-800-526-7736

Reference ID: 3612163

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Revised: April 2014